

# チャ

## カンザワハダニ

発生量は山城、丹波及び丹後で平年比やや多い。

## チャトゲコナジラミ

発生量は山城で例年比やや少なく、丹波、丹後で例年並。

## その他の病害虫

チュウゴクアミガサハゴロモの産卵痕が散見された。  
チャノコカクモンハマキ綴葉、幼虫が散見された。

### 令和8年 チャ害虫調査(カンザワハダニ・50葉調査)

調査日:山城:3/23、24 丹波・丹後:3/24

調査場所	カンザワハダニ						チャトゲコナジラミ						備考	
	寄生 葉率 (%)	100葉当たり			産卵 葉率 (%)	100葉当たり 卵数	発 生 ほ場率 (%)	寄生 葉率 (%)	発生程度別葉数			発生量 指数		発生ほ場 率(%)
		成虫数	幼虫数	合計					0	1	2			
(山城地域)														
宇治市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市3	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市4	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市5	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市6	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
城陽市1	0.0	0	0	0	0.0	0		18.0	41	7	2	11		
城陽市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		チュウゴクアミガサハゴロモの産卵痕を確認
宇治田原町1	2.0	2	0	2	0.0	0		4.0	48	2	0	2		
宇治田原町2	0.0	0	0	0	0.0	0		2.0	49	1	0	1		チュウゴクアミガサハゴロモの産卵痕を確認
宇治田原町3	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治田原町4	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
京田辺市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
京田辺市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
木津川市1	0.0	0	0	0	0.0	0		2.0	49	1	0	1		
木津川市2	0.0	0	0	0	0.0	0		16.0	42	8	0	8		チャノコカクモンハマキの幼虫を確認
和束町1	0.0	0	0	0	0.0	0		12.0	44	6	0	6		
和束町2	2.0	2	0	2	2.0	12		2.0	49	1	0	1		
和束町3	6.0	5	0	5	2.0	22		4.0	48	2	0	2		
和束町4	0.0	0	0	0	0.0	0		32.0	34	14	2	18		
和束町5	0.0	0	0	0	0.0	0		6.0	47	2	1	4		
和束町6	4.0	4	0	4	4.0	40		0.0	50	0	0	0		
平均値	0.6	0.6	0.0	0.6	0.4	3.4	27.3	4.5	47.8	2.0	0.2	2.5	45.5	
平年比	並	並	並	並	並	並	やや多	(やや少)	—	—	—	(並)	(やや少)	
前年値	1.4	1.6	0.6	2.2	1.0	21.2	45.5	6.7	—	—	—	3.5	59.1	
平年値(例年値)	0.6	0.7	0.3	1.0	0.3	5.5	19.1	(13.3)				(7.6)	(62.4)	
(丹波地域)														
福知山市1	8.0	10	0	10	8.0	130		0.0	50	0	0	0		
福知山市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
綾部市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
綾部市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
舞鶴市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
舞鶴市2	0.0	0	0	0	0.0	0		4.0	48	2	0	2		
平均値	1.3	1.7	0.0	1.7	1.3	21.7	16.7	0.7	49.7	0.3	0.0	0.3	16.7	
平年比	多	多	並	多	多	やや多	並	(並)	—	—	—	(やや少)	(並)	
前年値	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.7	—	—	—	1.8	66.7	
平年値	0.5	0.7	0.0	0.7	0.1	5.5	20.0	(6.8)				(3.6)	(45.8)	
(丹後地域)														
京丹後市1	0.0	0	0	0	0.0	0		16.0	42	8	0	8		
京丹後市2	0.0	0	0	0	0.0	0		20.0	40	10	0	10		チャノコカクモンハマキの幼虫を確認
京丹後市3	2.0	4	0	4	2.0	30		18.0	41	9	0	9		チャノコカクモンハマキの綴り葉を確認
京丹後市4	0.0	0	0	0	0.0	0		14.0	43	6	1	8		
平均値	0.5	1.0	0.0	1.0	0.5	7.5	25.0	17.0	41.5	8.3	0.3	8.8	100.0	
平年比	並	やや多	並	やや多	多	多	並	(並)	—	—	—	(並)	(並)	
前年値	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8.0	—	—	—	4.5	100.0	
平年値(例年値)	0.4	0.6	0.0	0.6	0.1	1.0	17.5	(17.7)				(9.7)	(81.3)	

カンザワハダニ(3月第5半旬)

地域	平年比	項目 \ 年次	R8	平年比	平年	R7	R6	R5	R4	R3	R2	H31	H30	H29	H28
山城	やや多	寄生葉率(%)	0.6	並	0.6	1.4	0.1	0.5	0.9	0.2	0.1	0.1	0.9	0.4	1.2
		成虫数(100葉当)	0.6	並	0.7	1.6	0.1	1.0	1.1	0.4	0.1	0.1	0.6	0.4	1.6
		幼虫数( " )	0.0	並	0.3	0.6	0.0	1.4	0.1	0.0	0.0	0.0	0.4	0.0	0.8
		合計( " )	0.6	並	1.0	2.2	0.1	2.4	1.2	0.4	0.1	0.1	1.0	0.4	2.5
		産卵葉率(%)	0.4	並	0.3	1.0	0.0	0.4	0.6	0.1	0.1	0.0	0.1	0.0	0.5
		卵数(100葉当)	3.4	並	5.5	21.2	0.0	10.8	8.9	4.4	0.4	0.0	0.9	0.0	8.0
		発生ほ場率(%)	27.3	やや多	19.1	45.5	4.5	22.7	22.7	9.1	4.8	4.5	27.3	13.6	36.4
丹波	やや多	寄生葉率(%)	1.3	多	0.5	0.0	0.7	1.3	0.0	0.3	1.0	0.3	0.3	0.0	1.0
		成虫数(100葉当)	1.7	多	0.7	0.0	1.3	1.3	0.0	0.3	1.3	1.3	0.3	0.0	1.0
		幼虫数( " )	0.0	並	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		合計( " )	1.7	多	0.7	0.0	1.3	1.3	0.0	0.3	1.3	1.3	0.3	0.0	1.0
		産卵葉率(%)	1.3	多	0.1	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0
		卵数(100葉当)	21.7	やや多	5.5	0.0	16.7	0.0	0.0	0.0	0.0	33.3	5.3	0.0	0.0
		発生ほ場率(%)	16.7	並	20.0	0.0	33.3	50.0	0.0	16.7	33.3	16.7	16.7	0.0	33.3
丹後	やや多	寄生葉率(%)	0.5	並	0.4	0.0	0.0	1.0	0.0	0.5	1.5	0.0	0.0	0.0	0.5
		成虫数(100葉当)	1.0	やや多	0.6	0.0	0.0	2.5	0.0	0.5	2.0	0.0	0.0	0.0	0.5
		幼虫数( " )	0.0	並	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		合計( " )	1.0	やや多	0.6	0.0	0.0	2.5	0.0	0.5	2.0	0.0	0.0	0.0	0.5
		産卵葉率(%)	0.5	多	0.1	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0
		卵数(100葉当)	7.5	多	1.0	0.0	0.0	0.0	0.0	10.0	0.0	0.0	0.0	0.0	0.0
		発生ほ場率(%)	25.0	並	17.5	0.0	0.0	50.0	0.0	25.0	75.0	0.0	0.0	0.0	25.0

チャトゲコナジラミ(3月第5半旬)

地 域	平年 (例年)比	項目 \ 年次	R8	平年 (例年)比	平年 (例年)	R7	R6	R5	R4	R3	R2	H31	H30	H29	H28
山城	(やや少)	寄生葉率(%)	4.5	(やや少)	(13.3)	6.7	3.3	4.7	7.0	14.2	34.3	19.55	16.73	—	—
		発生量指数	2.5	(並)	(7.6)	3.5	1.9	2.6	3.7	7.5	23.7	10.05	7.66	—	—
		発生ほ場率(%)	45.5	(やや少)	(62.4)	59.1	40.9	54.5	50.0	63.6	85.7	77.27	68.20	—	—
丹波	(並)	寄生葉率(%)	0.7	(並)	(6.8)	3.7	1.3	0.0	0.0	2.0	22.0	21.33	4.00	—	—
		発生量指数	0.3	(やや少)	(3.6)	1.8	0.7	0.0	0.0	1.0	12.0	11.00	2.00	—	—
		発生ほ場率(%)	16.7	(並)	(45.8)	66.7	33.3	0.0	0.0	16.7	100.0	100.00	50.00	—	—
丹後	(並)	寄生葉率(%)	17.0	(並)	(17.7)	8.0	13.5	6.5	3.0	12.5	47.0	44.50	6.50	—	—
		発生量指数	8.8	(並)	(9.7)	4.5	6.8	3.8	1.5	6.5	28.8	22.50	3.25	—	—
		発生ほ場率(%)	100.0	(並)	(81.3)	100.0	100.0	75.0	25.0	75.0	100.0	100.00	75.00	—	—